

New Horizons SDC Pluto Encounter

Calibrated Data Overview

During the migration to the Planetary Data System's (PDS) PDS4 data standards, this current description was adapted from the PDS3 dataset catalog file, including updates found in the KEM1 Encounter phase version, providing light edits to the text, format, flow, and to make the description to better conform to this PDS4 data collection.

Abstract

This data set contains Calibrated data taken by the New Horizons Student Dust Counter (SDC) instrument during the Pluto Encounter mission phase.

This data set contains SDC observations taken during the Approach (Jan-Jul, 2015), Encounter, Departure, and Transition mission sub-phases, including flyby observations taken on 14 July, 2015, and departure and calibration data through late October, 2016. This data set completes the Pluto mission phase deliveries for SDC.

Changes since prior versions include the final batch of Pluto mission phase data, downlinked between the end of January, 2016 and late in October, 2016, including a Stim calibration in July.

These data were migrated from the previously released PDS3 data set NH-A-SDC-3-PLUTO-V3.0.

Data Set Overview

This data set contains Raw data taken by the New Horizons Student Dust Counter (SDC) instrument during the Pluto Encounter mission phase. The closest approach to Pluto occurred on July 14, 2015, at approximately 11:50 UTC.

The mission of the SDC is to analyze the size and distribution of Interplanetary Dust Particles (IDPs) along the New Horizons trajectory to the Kuiper Belt. SDC comprises twelve thin, permanently polarized polyvinylidene fluoride (PVDF) plastic film sensors, with a combined area of about 0.1 m², mounted on the top surface of a support panel and normal to the spacecraft ram direction (flight velocity). In addition, there are two reference sensors, identical to the top surface sensors, mounted on the back side of the detector support panel and protected from any dust impacts, used to monitor background noise levels.

An impacting IDP causes a depolarization charge when it penetrates the PVDF film on one of the sensors. That charge is then measured by that sensor's electronics (channel); if the measurement is above a preset level, the instrument records and stores the event for later downlink. The level preset is adjusted based on in-flight Noise Floor Calibrations, and there are extensive autonomy rules adjusting SDC behavior, even turning channels off for up to thirty days at a time, to avoid overloading the storage system with noise.

SDC was designed to detect events for particles down to about one picogram at Pluto (see Bagenalet al. (2016)); that detection limit is lower than earlier in the mission where the

spacecraft velocity was higher. The SDC instrument has a temperature- and velocity-dependent calibration, first converting the raw measurement to charge, then converting charge to particle mass.

The common data product is a binary table of downlinked event data: time; sensor channel; magnitude; threshold magnitude. Associated data products are housekeeping data such as instrument temperatures for calibration and near-in-time spacecraft thruster events, which may induce false positives i.e. SDC events not caused by IDPs. The channels in the binary table for raw data are numbered from 0 to 13; the channel in the binary table for calibrated data are numbered from 1 to 14.

Some time between instrument delivery to the spacecraft and launch, the detector on one channel began exhibiting symptoms of degraded electrical contacts to the PVDF; data from that channel (channel number 10 in raw data; channel number 11 in calibrated data) are still processed but should be ignored.

During the Pluto Charon Encounter mission phase starting in January, 2015, there were several sub-phases: three Approach sub-phases, (AP1, AP2 and AP3); a CORE sequence for the Pluto flyby on 14 July, 2015 (Day Of Year 195), sometimes also referred to as NEP (Near-Encounter Phase); three Departure sub-phases (DP1, DP2, DP3). For this third and final SDC delivery for the Pluto mission phase, this data set includes all SDC data through the end of the Pluto mission phase in late-October, 2016, including all encounter data.

SDC was turned on throughout Approach. It was powered off then on periodically when needed for spacecraft power reasons or for trajectory correction maneuvers, and the channels were turned off and on around the start of DSN tracks that contained tweakups, as per normal operations. The thresholds were updated and set higher for Pluto - making SDC less sensitive to dust hits - on January 1st, 2015 (DOY 001), then set back to Cruise threshold settings on July 30th (DOY 211). The threshold used for each dust hit is recorded in extension 1 of the calibrated data products. A Stimulus calibration was performed on July 08, 2016.

During the five days before and after the Pluto flyby, SDC detected one probable dust hit event. The event happened on July 11, three days prior to Pluto closest approach, at a distance of ~3000 Pluto radii. Refer to figure 6 of Bagenal et al. (2016) for more details.

Every observation provided in this data set was taken as a part of a particular sequence. For this data set, these sequences can be found in the SDC document collection under PDS4 LID `urn:nasa:pds:nh_documents:sd:seq_sdc_pluto`. Please note that some sequences provided may have zero corresponding observations.

Version History

Each subsection below details the major changes between the prior versions of this data set, listing the newest versions before older versions.

[PDS4 v1.0 \(migration from PDS3 V3.0\)](#)

This data collection was migrated from Planetary Data System's (PDS) PDS3 archive standards to the PDS4 archive standards, which involved changing the PDS formatted product labels. The products themselves have remained unchanged. The major changes from the PDS3 V3.0 data set are:

- the calibration files, documents, and data products were reorganized into separate collections of calibration files, documents, and data products, instead of being in a single package as it was in prior PDS3 data set versions.
- the geometry keyword values found within the PDS4 labels were calculated using the most recent SPICE kernels available at label creation. Note that the FITS headers have not been updated and their geometry keyword values therefore remain unchanged.
- the PDS4 data labels were produced using the PDS3 data labels and/or FITS headers, and so any fixes and/or updates to the PDS3 label pipeline as found in future mission phases may not have been implemented here.

[PDS3 V3.0 \(NH-P-SDC-3-PLUTO-V3.0\)](#)

This is VERSION 3.0 of this data set. This P3 Pluto Encounter dataset release provides updates to the Pluto dataset between P2 (data on the ground by 1/31/2016) and P3 (data on the ground by the end of the Pluto mission phase in late October, 2016); this completes the SDC Pluto data set. Since the P2 delivery data, SDC has stayed on and taken data continuously, however, see the comment regarding tweakups in the V2.0 updates below. A Stim calibration was performed on 07/08/2016.

Also, updates were made to the documentation and catalog files, primarily to implement suggestions from the V2.0 peer review. A new table of SDC Ram (velocity) ancillary data has been provided, and the SDC on/off and Stim tables have been extended in time to cover the new data.

PDS Citation Information: Horányi M., NEW HORIZONS Calibrated SDC PLUTO ENCOUNTER V3.0, NH-P-SDC-3-PLUTO-V3.0, NASA Planetary Data System, 2018.

[PDS3 V2.0 \(NH-P-SDC-3-PLUTO-V2.0\)](#)

This is VERSION 2.0 of this data set. This P2 Pluto Encounter dataset release provides updates to the Pluto dataset between P1 (data on the ground by 7/31/2015) and P2 (data on the ground by 1/31/2016). All liens from the initial Pluto delivery have also now been resolved. For SDC, most of the Pluto Encounter data was downlinked in the 15229 load in August 2015. Since then, SDC has stayed on and taken data continuously, due to power restrictions it has frequently been turned off anytime there was a downlink of SSR Side 2 data. The channels are also turned off and on around the start of certain DSN tracks, about every 3 days (these tracks are known as tweakups).

PDS Citation Information: Horányi M., NEW HORIZONS Calibrated SDC PLUTO ENCOUNTER V2.0, NH-P-SDC-3-PLUTO-V2.0, NASA Planetary Data System, 2017.

PDS3 V1.0 (NH-P-SDC-3-PLUTO-V1.0)

This is VERSION 1.0 of this data set. For this first SDC delivery for the Pluto mission phase, this data set includes all SDC data through late-July, including all encounter data. Liens were never resolved for this data set version but will be in the next version.

PDS Citation Information: Horányi M., NEW HORIZONS Calibrated SDC PLUTO ENCOUNTER V1.0, NH-P-SDC-3-PLUTO-V1.0, NASA Planetary Data System, 2016.

General statement about data set versions after V1.0

The pipeline (see Processing below) was re-run on these data for each version since the first (V1.0). A pipeline rerun usually changes the FITS headers but not the FITS data of raw data sets. In some cases, calibrated FITS data may change because the calculated geometry of an observation has changed. See data set version-specific sections above for significant exceptions to this general statement, i.e., changes to pipeline processing, calibration processing, and data delivered.

An all-instrument Calibration Campaign occurred in July 2016. For most instruments, calibrations were updated as of April 2017 which changed the data in the calibrated data sets. Calibration changes are described in the data set version-specific sections.

Note that even if this is not a calibrated data set, calibration changes are listed as the data will have been re-run and there will be updates to the calibration files, to the documentation and to the steps required to calibrate the data.

Also note that file names may change between versions if start/stop times are updated when additional data are downlinked.

Processing

The data in this data set were created by a software data processing pipeline on the Science Operations Center (SOC) at the Southwest Research Institute (SwRI), Department of Space Operations. This SOC pipeline assembled data as FITS files from raw telemetry packets sent down by the spacecraft and populated the data labels with housekeeping and engineering values, and computed geometry parameters using SPICE kernels. The pipeline did not resample the data.

SDC data calibration is a two-step process: raw data numbers from a particle impact are converted to a charge, and the charge is converted to a particle mass via the ground calibrations obtained at a dust acceleration facility. Refer to the provided documentation for more information. The latest calibration procedure is described in James et al. (2010).

Data

The observations in this data set are stored in data files using standard Flexible Image Transport System (FITS) format. Each FITS file has a corresponding detached PDS label file, named according to a common convention. The FITS files may have image and/or table extensions.

- start_date_time
- stop_date_time
- start_clock_count
- stop_clock_count

Also note that a dust hit may be recorded in more than one data product. The product name reflects the MET when data was transferred from instrument memory. More than one transfer may have been requested for any given interval from start_date_time to stop_date_time. When processing the data, check these intervals carefully. Obvious cases of duplicate dust hit data spanning long time intervals have already been removed from this dataset.

Application ID (ApID)

Here is a summary of the types of files generated by each ApID (N.B. ApIDs are case-insensitive) along with the instrument designator that go with each ApID:

ApIDs	Data product description/Prefix(es)
0x700	SDC Science Data/SDC

There are other ApIDs that contain housekeeping values and other values. See the SOC Instrument ICD for more details: `urn:nasa:pds:nh_documents:mission:soc_inst_icd`

Please note that not all ApIDs may be found in this data set.

Instrument description

Refer to the following files for a description of this instrument:

- New Horizon SDC instrument overview: `urn:nasa:pds:nh_documents:sdcsdc_inst_overview`
- SDC Space Science Review (SSR) paper: `urn:nasa:pds:nh_documents:sdcsdc_ssr`
- SOC Instrument ICD: `urn:nasa:pds:nh_documents:mission:soc_inst_icd`
- SDC SPICE Instrument Kernel: `urn:nasa:pds:nh_documents:sdcsnh_sdc_ti`

Other sources of information useful in interpreting these Data

Refer to the following files for more information about these data:

- NH Mission Trajectory Table: `urn:nasa:pds:nh_documents:mission:nh_mission_trajectory`
- Field of View Illustration: `urn:nasa:pds:nh_documents:mission:nh_fov`
- SDC SPICE Instrument Kernel: `urn:nasa:pds:nh_documents:sdcsnh_sdc_ti`
- SDC Summary Student Dust Counter (SDC) boresight direction: `urn:nasa:pds:nh_documents:sdcsdcrsam`

Visit Description, Visit Number, and Target in the Data Labels

The observation sequences were defined in Science Activity Planning (SAP) documents and grouped by Visit Description and Visit Number. The SAPs are spreadsheets with one Visit Description & Number per row. A nominal target is also included on each row and included in

the data labels but does not always match with the target name field's value in the data labels. In some cases, the target was designated as right_ascension_angle, declination_angle pointing values in the form "right_ascension_angle, declination_angle =123.45,-12.34" indicating Right Ascension and Declination, in degrees, of the target from the spacecraft in the Earth Equatorial J2000 inertial reference frame. This indicates that either the target was a star, or the target's ephemeris was not loaded into the spacecraft's attitude and control system which in turn meant the spacecraft could not be pointed at the target by a body identifier and an inertial pointing value had to be specified as Right Ascension and Declination values. PDS-SBN practices do not allow putting a value like right_ascension_angle, declination_angle =... in the PDS target name keyword's value. In those cases, the PDS target purpose value is set calibration. Target name may be None for a few observations in this data set; typically, that means the observation is a functional test so None is an appropriate entry for those targets, but the PDS user should also check the nh:observation_description and nh:sequence_id keywords in the PDS label, plus the provided sequence list (PDS4 LID `urn:nasa:pds:nh_documents:sdcs:seq_sdc_pluto`) to assess the possibility that there was an intended target. These two keywords are especially useful for star targets as often stars are used as part of instrument calibrations and are included as part of the sequencing description which is captured in these keywords.

For SDC the target name is always Dust.

Ancillary Data

The geometry items included in the data labels were computed using the SPICE kernels archived in the New Horizons SPICE data set, NH-J/P/SS-SPICE-6-V1.0, <https://doi.org/10.17189/1520109>.

Every observation provided in this data set was taken as a part of a particular sequence. A list of these sequences has been provided within the NH SDC document collection (PDS4 LID `urn:nasa:pds:nh_documents:sdcs`) within the PDS, one file for each mission phase. The sequence identifier and description are included in the PDS label for every observation.

N.B. While every observation has an associated sequence, every sequence may not have associated observations. Some sequences may have failed to execute due to spacecraft events (e.g., safing). No attempt has been made during the preparation of this data set to identify such empty sequences.

Time

There are several time systems, or units, in use in this dataset: New Horizons spacecraft MET (Mission Event Time or Mission Elapsed Time), UTC (Coordinated Universal Time), and TDB (Barycentric Dynamical Time).

This section will give a summary description of the relationship between these time systems. For a complete explanation of these time systems the reader is referred to the documentation distributed with the Navigation and Ancillary Information Facility (NAIF) SPICE toolkit from the PDS NAIF node, (see <http://naif.jpl.nasa.gov/>).

The most common time unit associated with the data is the spacecraft MET. MET is a 32-bit counter on the New Horizons spacecraft that runs at a rate of about one increment per second starting from a value of zero at “19.January, 2006 18:08:02 UTC” or “JD2453755.256337 TDB.”

The leapsecond adjustment ($\Delta ET = ET - UTC$) was 65.184s at NH launch, and the first four additional leapseconds occurred at the ends of 12/2009, 06/2012, 06/2015, and 12/2016. Refer to the NH SPICE data set, NH-J/P/SS-SPICE-6-V1.0, <https://doi.org/10.17189/1520109>, and the SPICE toolkit documentation, for more details about leapseconds.

The data labels for any given product in this dataset usually contain at least one pair of common UTC and MET representations of the time at the middle of the observation. Other portions of the products, for example tables of data taken over periods of up to a day or more, will only have the MET time associated with a given row of the table.

For the data user's use in interpreting these times, a reasonable approximation ($\pm 1s$) of the conversion between Julian Day (TDB) and MET is as follows:

$$JD_{TDB} = 2453755.256337 + (MET / 86399.9998693)$$

For more accurate calculations the reader is referred to the NAIF/SPICE documentation as mentioned above.

Reference Frame

Geometric Parameter Reference Frame

Earth Mean Equator and Vernal Equinox of J2000 (EMEJ2000) is the inertial reference frame used to specify observational geometry items provided in the data labels. Geometric parameters are based on best available SPICE data at time of data creation.

Epoch of Geometric Parameters

All geometric parameters provided in the data labels were computed at the epoch midway between the start_date_time and stop_date_time label fields.

Software

The observations in this data set are in standard FITS format with PDS labels and can be viewed by a number of PDS-provided and commercial programs. For this reason, no special software is provided with this data set.

Confidence Level Overview

During the processing of the data in preparation for delivery with this volume, the packet data associated with each observation were used only if they passed a rigorous verification process including standard checksums.

In addition, raw (CODMAC Level 2) observation data for which adequate contemporary housekeeping and other ancillary data are not available may not be reduced to calibrated

(CODMAC Level 3) data. This issue is raised here to explain why some data products in the raw data set may not have corresponding data products in the calibrated data set.

Data coverage and quality

Every observation provided in this data set was taken as a part of a particular sequence. For this data set, these sequences can be found in the SDC document collection under PDS4 LID `urn:nasa:pds:nh_documents:sdsc:seq_sdc_pluto`. Please note that some sequences provided may have zero corresponding observations.

Refer to the Confidence Level Overview section above for a summary of steps taken to assure data quality.

For SDC, the stimulus calibration activity is known to generate false positive events in the science data. This data set includes a table, found within the SDC document collection under PDS4 LID `urn:nasa:pds:nh_documents:sdsc:sdsc_stim`, that lists time periods when stimulus calibrations were active (several times during Launch and Jupiter mission phases, and about half an hour per year during Annual CheckOuts (ACO) in the Pluto Cruise mission phase. Eventually, the Science Operations Center (SOC) operational pipeline may be enhanced to filter individual events that occur near stimulus events.

Caveat about target name in PDS labels and observational

The downlink team on New Horizons has created an automated system to take various uplink products, decode things like Chebyshev polynomials in command sequences representing celestial body ephemerides for use on the spacecraft to control pointing, and infer from those data what the most likely intended target was at any time during the mission. This works well during flyby encounters and less so during cruise phases and hibernation.

The user of these PDS data needs to be cautious when using the target name and other target-related parameters stored in this data set. This is less an issue for the plasma and particle instruments, more so for pointed instruments. To this end, the heliocentric ephemeris of the spacecraft, the spacecraft-relative ephemeris of the inferred target, and the inertial attitude of the instrument reference frame are provided with all data, in the J2000 inertial reference frame, so the user can check where that target is in the Field Of View (FOV) of the instrument.

Finally, note that, within the FITS headers of the data products, the sequence tables, and other NH Project-internal documents used in this data set, informal names are often used for targets instead of the canonical names used within the PDS labels. For example, during the Pluto mission phase, instead of the target name '15810 ARAWN (1994 JR1)' there might be found any of the following: 1994JR1; 1994 JR1; JR1. However, within the context of this data set, these project abbreviations are not ambiguous (e.g. there is only one NH target with 'JR1' in its name), so there has been, and will be, no attempt to expand such abbreviations where they occur outside formal PDS keyword values.

Contact Information

For any questions regarding the data format of the archive, contact the New Horizons SDC Principal Investigator:

Mihály Horányi
Laboratory for Atmospheric and Space Physics
University of Colorado
Boulder, CO 80302-3092
USA

Reference List

Bagenal, F., M. Horányi D.J. McComas, R.L. McNutt Jr, H.A. Elliott, M.E. Hill, L.E. Brown, P.A. Delamere, P. Kollmann, S.M. Krimigis, M. Kusterer, C.M. Lisse, D.G. Mitchell, M. Piquette, A.R. Poppe, D.F. Strobel, J.R. Szalay, P. Valek, J. Vandegriff, S. Weidner, E.J. Zirnstein, S.A. Stern, K. Ennico, C.B. Olkin, H.A. Weaver, L.A. Young, and New Horizons Science Team, Pluto's interaction with its space environment: Solar wind, energetic particles & dust, Science, Volume 351, Issue 6279, 18 Mar 2016. <https://doi.org/10.1126/science.aad9045>

Further Reading

James, D., V. Hoxie, and M. Horanyi, Polyvinylidene fluoride dust detector response to particle impacts, Review of Scientific Instruments, Volume 81, Issue 3, id. 034501-034501-8, 2010. <https://doi.org/10.1063/1.3340880>

Steffl, A.J., J. Peterson, B. Carcich, L. Nguyen, and S.A. Stern, NEW HORIZONS SPICE KERNELS, V1.0, NH-J/P/SS-SPICE-6-V1.0, NASA Planetary Data System, 2007. <https://doi.org/10.17189/1520109>